

**ORIGINAL RESEARCH ARTICLE**
**Immediate onset signatures of autoimmune diseases after vaccination**
**Darrell O. Ricke\***

Department of Research, Molecular BioInsights, Winchester, MA, 01890, USA

**Supplementary File**
**Table S1. Autoimmune adverse events by vaccine type for COVID-19, FLU, HEP, HPV, PNC, and VARZOS**

Adverse event	COVID-19	FLU	HEP	HPV	PNC	VARZOS
Common adverse events: Headache, fatigue, pyrexia (fever), chills, and pain	898,859 240,413 26.75%	92,682 34,067 36.76%	36,222 13,914 38.41%	21,437 9,107 42.48%	23,491 10,859 46.23%	69,901 23,118 33.07%
Acute disseminated encephalomyelitis	280 90 0.01%	205 85 0.09%	75 31 0.09%	79 30 0.14%*	34 13 0.06%	17 7 0.01%
Alopecia	3,991 1,621 0.18%	156 62 0.07%	390 146 0.40%*	653 369 1.72%*	37 16 0.07%	102 60 0.09%
Arthritis	4,199 1,759 0.20%	501 191 0.21%	655 264 0.73%*	223 64 0.30%	51 29 0.12%	188 122 0.18%
Autoimmune disorder	1,709 770 0.09%	142 54 0.06%	131 66 0.18%*	336 171 0.80%*	- - -	84 55 0.08%
Autoimmune hepatitis	439 162 0.02%	14 6 0.01%	49 36 0.10%	22 9 0.04%	- - -	- - -
Autoimmune thyroiditis	770 299 0.03%	- - -	84 17 0.05%	117 31 0.15%*	- - -	- - -
Bell's palsy	7,331 2,129 0.24%*	205 37 0.04%	- - -	- - -	- - -	146 82 0.12%*
Chronic fatigue syndrome	905 451 0.05%	30 12 0.01%	89 49 0.14%*	276 178 0.83%*	- - -	- - -
Complex regional pain syndrome	168 96 0.01%	- - -	- - -	150 64 0.30%*	- - -	- - -

*(Cont'd...)*

Table S1. (Continued)

Adverse event	COVID-19	FLU	HEP	HPV	PNC	VARZOS
Crohn's disease	799 382 0.04%	19 6 0.00%	62 39 0.11%*	80 36 0.17%*	- - -	9 6 0.01%
Diabetes mellitus	1,159 485 0.05%	116 46 0.05%	176 47 0.13%	44 19 0.09%	45 24 0.10%	17 11 0.02%
Eczema	2,117 872 0.10%	174 68 0.07%	244 96 0.27%*	107 44 0.21%*	306 165 0.70%*	86 41 0.06%
Guillain-Barré syndrome (GBS)	3,451 1,278 0.14%	3,045 869 0.94%*	439 106 0.29%*	276 126 0.59%*	172 47 0.20%	396 201 0.29%*
Henoch-Schonlein purpura	259 85 0.01%	153 26 0.03%	109 12 0.03%	107 19 0.09%	65 18 0.08%	- - -
Immune thrombocytopenia	1,667 581 0.06%	39 11 0.01%	33 8 0.02%	- - -	64 13 0.06%	- - -
Juvenile idiopathic arthritis	56 17 0.00%	- - -	- - -	78 43 0.20%*	11 5 0.02%	- - -
Kawasaki's disease	57 16 0.00%	39 11 0.01%	100 24 0.07%	- - -	263 101 0.43%*	- - -
Lyme disease	517 198 0.02%	- - -	- - -	54 28 0.13%*	- - -	15 15 0.02%
Multiple sclerosis	1230 535 0.06%	151 56 0.06%	885 427 1.18%*	306 186 0.87%*	- - -	32 26 0.04%
Multiple sclerosis relapse	842 365 0.04%	28 12 0.01%	37 16 0.04%	13 6 0.03%	- - -	21 16 0.02%
Myasthenia gravis	624 233 0.03%	30 8 0.01%	- - -	36 18 0.08%	- - -	- - -
Myositis	760 337 0.04%	183 70 0.08%	115 34 0.09%	52 15 0.07%	22 7 0.03%	21 9 0.01%
Narcolepsy	158 71 0.01%	170 93 0.10%	15 7 0.02%	116 73 0.34%*	- - -	- - -

(Cont'd...)

Table S1. (Continued)

Adverse event	COVID-19	FLU	HEP	HPV	PNC	VARZOS
Neutropenia	494	53	85	29	106	-
	148	17	7	7	52	-
	0.02%	0.02%	0.02%	0.03%	0.22%*	-
Optic neuritis	831	181	335	155	-	31
	271	8	128	51	-	17
	0.03%	0.01%	0.35%*	0.24%*	-	0.02%
Polymyalgia rheumatica	1,994	186	28	-	17	146
	691	75	12	-	11	75
	0.08%	0.08%	0.03%	-	0.05%	0.11%
Postural orthostatic tachycardia syndrome (POTS)	838	35	13	665	-	-
	334	6	5	409	-	-
	0.04%	0.01%	0.01%	1.91%*	-	-
Raynaud's phenomenon	586	20	44	83	-	-
	218	10	13	42	-	-
	0.02%	0.01%	0.04%	0.20%*	-	-
Restless legs syndrome	742	-	29	48	-	21
	399	-	16	27	-	11
	0.04%	-	0.04%	0.20%*	-	0.02%
Rheumatoid arthritis	3,411	195	337	113	25	161
	1,552	92	155	47	15	107
	0.17%*	0.10%	0.43%*	0.22%*	0.06%	0.15%*
Sjogren's syndrome	354	-	45	37	-	-
	132	-	32	23	-	-
	0.02%	-	0.09%*	0.11%*	-	-
Systemic lupus erythematosus	802	112	242	256	-	38
	377	61	112	149	-	24
	0.04%	0.07%	0.31%*	0.70%*	-	0.03%
Type 1 diabetes mellitus	647	65	68	108	31	-
	208	11	23	34	7	-
	0.02%	0.01%	0.06%	0.16%*	0.03%	-
Uveitis	726	42	125	72	-	43
	273	25	96	45	-	23
	0.03%	0.03%	0.27%*	0.21%*	-	0.03%
Vasculitis	1,085	213	279	62	61	37
	377	90	117	29	27	21
	0.04%	0.10%	0.32%*	0.14%*	0.12%*	0.03%

Notes: First row: total number of reports; second row: number of reports on day 0; third row: percentage of the number of reports on day 0 over total number of reports for the relevant vaccine type. \*The percentage that is high compared to other percentages for the same adverse event. VAERS data include adverse event reports from 1990 to May 19, 2023. FLU: Influenza, HEP: Hepatitis A and B, HPV: Human papillomavirus, PNC: Pneumococcal, and VARZOS: Varicella-zoster.

**Table S2. Adverse event enrichment estimates over background expected for onset within the first 24 h**

Adverse event	Vaccine	Background per 100,000/year	b <sup>x</sup>	AE <sub>0</sub> reports	AE <sub>0</sub> per 100,000/year	V <sub>0</sub> ( $r_o = 1/20$ )	V <sub>0</sub> ( $r_o = 1/40$ )
Alopecia	COVID-19	20.75 <sup>[1]</sup>	0.057	1,621	288	15.7	31.5
Bell's palsy	COVID-19	23.8 <sup>[2]</sup>	0.065	2,129	378	20.6	41.3
Chronic fatigue syndrome	COVID-19	13.16 <sup>[3]</sup>	0.036	451	80	4.3	8.7
Complex regional pain syndrome	COVID-19	5.46 <sup>[4]</sup>	0.015	96	17	0.9	1.9
Guillain-Barré syndrome	COVID-19	2.1 <sup>[2]</sup>	5.75E-03	1,278	227	12.4	24.8
Henoch-Schonlein purpura	COVID-19	1.93 <sup>[5]</sup>	5.29E-03	85	15	0.8	1.6
Immune thrombocytopenia	COVID-19	21.7 <sup>[2]</sup>	5.95E-02	581	103	5.6	11.2
Myositis	COVID-19	22	0.060	337	60	3.2	6.5
Multiple sclerosis	COVID-19	2.1 <sup>[6]</sup>	0.006	535	95	5.2	10.4
Narcolepsy	COVID-19	1.1 <sup>[2]</sup>	3.01E-03	71	13	0.7	1.4
Optic neuritis	COVID-19	3.9 <sup>[7]</sup>	0.011	271	48	2.6	5.3
POTS	COVID-19	10.1 <sup>[8]</sup>	0.028	334	59	3.2	6.5
Rheumatoid arthritis	COVID-19	41 <sup>[9]</sup>	0.112	1,552	275	15.0	30.1
Systemic lupus erythematosus	COVID-19	5.3 <sup>[2]</sup>	0.015	377	67	3.7	7.3
Type 1 diabetes mellitus	COVID-19	15 <sup>[10]</sup>	0.041	208	37	2.0	4.0
Guillain-Barré syndrome	Influenza	2.1	0.01	84	17	1.0	1.9
Alopecia	Hepatitis	30.75	0.08	146	53	2.8	5.8
Chronic fatigue syndrome	Hepatitis	13.16	3.61E-02	49	18	0.9	1.9
Guillain-Barré syndrome	Hepatitis	2.1	5.75E-03	106	39	2.1	4.2
Multiple sclerosis	Hepatitis	2.1	0.01	427	156	8.5	17.1
Optic neuritis	Hepatitis	3.9	1.07E-02	128	47	2.5	5.1
Systemic lupus erythematosus	Hepatitis	5.3	0.01	112	41	2.2	4.5
Vasculitis	Hepatitis	0.0034	9.32E-06	117	43	2.3	4.7
Alopecia	HPV	20.75	0.06	369	101	5.5	11.0
Chronic fatigue syndrome	HPV	13.16	3.61E-02	178	49	2.6	5.3
Complex regional pain syndrome	HPV	5.46	1.50E-02	64	18	0.9	1.9
Guillain-Barré syndrome	HPV	2.1	5.75E-03	126	34	1.9	3.8
Multiple sclerosis	HPV	2.1	0.01	186	51	2.8	5.6
Narcolepsy	HPV	1.1	3.01E-03	73	20	1.1	2.2
Optic neuritis	HPV	3.9	1.07E-02	51	14	0.8	1.5
POTS	HPV	10.1	2.77E-02	334	91	5.0	10.0
Systemic lupus erythematosus	HPV	5.3	0.01	149	41	2.2	4.5
Guillain-Barré syndrome	PNC	2.1	5.75E-03	47	33	1.8	3.6
Kawasaki disease	PNC	18.1 <sup>[11]</sup>	4.96E-02	101	71	3.8	7.7
Neutropenia	PNC	1 <sup>[12]</sup>	2.74E-03	52	36	2.0	4.0
Bell's palsy	VARZOS	23.8	0.065	82	67	3.6	7.3
Guillain-Barré syndrome	VARZOS	2.1	5.75E-03	201	164	9.0	18.0
Rheumatoid arthritis	VARZOS	41	0.112	107	87	4.7	9.4

Notes: Immunized population estimates used: COVID-19: 617,135,227 doses with three shots per person; Influenza: 175 million (averaged 2021, 2022, and 2023 reports); Hepatitis: 30% of the U.S. adult population of 333,287,557; Human papillomavirus (HPV): 40% of the U.S. adult population of 333,287,557, Pneumococcal (PNC): 70% of older adults of 74.6 million; Varicella-zoster (VARZOS): 60% of older adults of 74.6 million. AE<sub>0</sub>: adverse event reported on day 0; B<sup>x</sup>: background adverse events; V<sub>0</sub>: Vaccine-associated adverse events reported on day 0 (within 24 hours) of immunization.

**Table S3. Autoimmune adverse events induced by other vaccines**

Adverse event	6VAX-F	ANTH	HIBV	IPV	LYME	MENB	MNQ	PPV	RV	TYP
Common adverse events	1,683	4,792	20,668	11,990	1,788	10,055	9,356	35,144	6,400	4,944
Arthritis	-	104	52	-	-	-	-	117	-	-
	-	50	19	-	-	-	-	71	-	-
	-	1.04%*	0.09%	-	-	-	-	0.20%*	-	-
Guillain-Barré syndrome	-	-	49	81	-	23	134	207	-	77
	-	-	19	18	-	12	14	71	-	19
	-	-	0.09%	0.15%*	-	0.17%	0.15%*	0.20%*	-	0.38%*
Henoch-Schonlein purpura	-	-	29	41	-	26	-	22	-	-
	-	-	13	12	-	8	-	10	-	-
	-	-	0.06%	0.10%	-	0.11%	-	0.03%	-	-
Kawasaki disease	65	-	124	-	-	88	-	14	181	-
	30	-	47	-	-	50	-	12	53	-
	1.78%*	-	0.23%	-	-	0.69%*	-	0.03%	0.83%*	-
Neutropenia	33	-	44	32	-	34	-	25	54	-
	18	-	20	12	-	12	-	18	21	-
	1.07%*	-	0.10%	0.10%	-	0.17%	-	0.05%	0.33%*	-
Rheumatoid arthritis	-	28	-	-	84	-	-	38	-	-
	-	7	-	-	34	-	-	24	-	-
	-	0.15%	-	-	1.90%*	-	-	0.07%	-	-
Vasculitis	-	-	45	32	-	-	-	39	-	16
	-	-	21	16	-	-	-	17	-	5
	-	-	0.10%	0.13%	-	-	-	0.05%	-	0.10%

Notes: First row: Total number of reports; second row: Number of reports on day 0; third row: Percentage of the number of reports on day 0 over total number of reports for the relevant vaccine type. \*The percentage that is high compared to other percentages for the same adverse event. VAERS data include adverse event reports from 1990 to May 19, 2023. 6-VAX-F: Diphtheria and tetanus toxoids and acellular pertussis absorbed+inactivated poliovirus+hepatitis B+Haemophilus B conjugate vaccine; ANTH: Anthrax vaccine; HIBV: *Haemophilus influenzae* type B vaccine; IPV: Inactivated poliovirus vaccine; LYME: Lyme disease vaccine; MENB: Meningococcal group B, rDNA absorbed vaccine; MNQ: Meningococcal conjugate vaccine; PPV: Pneumococcal, polyvalent vaccine; RV: Rotavirus vaccine; TYP: Typhoid vaccine.

## References

1. Mirzoyev SA, Schrum AG, Davis MDP, *et al.*, 2014, Lifetime incidence risk of alopecia areata estimated at 2.1% by Rochester epidemiology project, 1990–2009. *J Invest Dermatol*, 134: 1141–1142.  
<https://doi.org/10.1038/jid.2013.464>
2. Willame C, Dodd C, van der Aa L, *et al.*, 2021, Incidence rates of autoimmune diseases in European healthcare databases: A contribution of the ADVANCE project. *Drug Saf*, 44: 383–395.  
<https://doi.org/10.1007/s40264-020-01031-1>
3. Vincent A, Brimmer DJ, Whipple MO, *et al.*, 2012, Prevalence, incidence, and classification of chronic fatigue syndrome in Olmsted County, Minnesota, as estimated using the Rochester epidemiology project. *Mayo Clin Proc*, 87: 1145–1152.  
<https://doi.org/10.1016/j.mayocp.2012.08.015>
4. Sandroni P, Benrud-Larson LM, McClelland RL, *et al.*, 2003, Complex regional pain syndrome Type I: Incidence and prevalence in Olmsted county, a population-based study. *Pain*, 103: 199–207.  
[https://doi.org/10.1016/S0304-3959\(03\)00065-4](https://doi.org/10.1016/S0304-3959(03)00065-4)
5. Tracy A, Subramanian A, Adderley NJ, *et al.*, 2019, Cardiovascular, thromboembolic and renal outcomes in IgA vasculitis (Henoch-Schönlein purpura): A retrospective cohort study using routinely collected primary care data. *Ann Rheum Dis*, 78: 261.  
<https://doi.org/10.1136/annrheumdis-2018-214142>
6. Walton C, King R, Baneke P, 2020, Rising prevalence of multiple sclerosis worldwide: Insights from the Atlas of MS, third edition. *Mult Scler*, 26: 1816–1821.  
<https://doi.org/10.1177/1352458520970841>
7. Hassan MB, Stern C, Flanagan EP, *et al.*, 2020, Population-based incidence of optic neuritis in the era of aquaporin-4

- and myelin oligodendrocyte glycoprotein antibodies. *Am J Ophthalmol*, 220: 110–114.  
<https://doi.org/10.1016/j.ajo.2020.07.014>
8. AbdelRazek M, Low P, Rocca W, et al., 2019, Epidemiology of postural tachycardia syndrome (S18.005). *Neurology*, 92: S18.005.
9. Myasoedova E, Davis J, Matteson EL, et al., 2020, Is the epidemiology of rheumatoid arthritis changing? Results from a population-based incidence study, 1985–2014. *Ann Rheum Dis*, 79: 440.  
<https://doi.org/10.1136/annrheumdis-2019-216694>
10. Mobasseri M, Shirmohammadi M, Amiri T, et al., 2020, Prevalence and incidence of Type 1 diabetes in the world: A systematic review and meta-analysis. *Health Promot Perspect*, 10: 98–115.  
<https://doi.org/10.34172/hpp.2020.18>
11. Elakabawi K, Lin J, Jiao F, et al., 2020, Kawasaki disease: Global burden and genetic background. *Cardiol Res*, 11: 9–14.  
<https://doi.org/10.14740/cr993>
12. Jinca C, Serban M, Ursu E, et al., 2021, Primary autoimmune neutropenia of infancy and childhood in a cohort of patients from western Romania. *Exp Ther Med*, 21: 280.  
<https://doi.org/10.3892/etm.2021.9711>